

The EEL-1 ubiquitin ligase promotes DNA damage-induced germ cell apoptosis in *C. elegans*

AJ Ross^{1,2}, M Li^{1,2}, B Yu¹, MX Gao¹ and WB Derry^{*1,2}

E3 ubiquitin ligases target a growing number of pro- and anti-apoptotic proteins, including tumour suppressor p53, caspases, and the Bcl-2 family. The core apoptosis pathway is well conserved between mammals and *Caenorhabditis elegans*, but the extent to which ubiquitin ligases regulate apoptotic cell death is not known. To investigate the role of E3 ligases in apoptosis, we inhibited 108 of the 165 predicted E3 ubiquitin ligase genes by RNA interference and quantified apoptosis in the *C. elegans* germline after genotoxic stress. From this screen, we identified the homologous to E6-associated protein C terminus-domain E3 ligase EEL-1 as a positive regulator of apoptosis. Intriguingly, the human homologue of EEL-1, Huwe1/ARF-BP1/Mule/HectH9, has been reported to possess both pro- and anti-apoptotic functions through its ability to stimulate Mcl-1 and p53 degradation, respectively. Here, we demonstrate that *eel-1* is required to promote DNA damage-induced germ cell apoptosis, but does not have a role in physiological germ cell apoptosis or developmental apoptosis in somatic tissue. Furthermore, *eel-1* acts in parallel to the p53-like gene *cep-1* and intersects the core apoptosis pathway upstream of the Bcl-2/Mcl-1 orthologue *ced-9*. Although *eel-1* mutants exhibit hypersensitivity to genotoxic stress they do not appear to be defective in DNA repair, suggesting a distinct role for EEL-1 in promoting damage-induced apoptosis in the germline.

Cell Death and Differentiation (2011) 18, 1140–1149; doi:10.1038/cdd.2010.180; published online 14 January 2011

E3 ubiquitin ligases act at the terminal step of an enzyme cascade that covalently attaches ubiquitin molecules to specific protein substrates. The process of ubiquitination involves either the attachment of a single ubiquitin moiety (monoubiquitination) or several ubiquitins linked in a chain (polyubiquitination) to a target protein. The E3 ligases comprise a large group of proteins subdivided by their domain architecture into the HECT (homologous to E6-associated protein C terminus), RING (really interesting new gene), and U-box (a modified RING motif) families.¹ Ubiquitin-based modifications generated by members of these E3 ligase families regulate myriads of biological processes, including cell-cycle progression, differentiation, DNA repair, and programmed cell death (PCD or apoptosis).¹ Deregulated expression of E3 ligases can result in a number of pathological conditions such as neurodegeneration and cancer. For example, endogenous levels of tumour suppressor p53 are under strict regulation by the RING-finger E3 ubiquitin ligase Mdm2 (murine double minute clone 2 oncoprotein), which targets p53 for ubiquitin-mediated proteolysis under normal conditions.² In response to stresses such as DNA damage, p53 and Mdm2 become phosphorylated and their interaction is inhibited, which leads to stabilization and activation of p53. In addition to Mdm2, other classes of E3 ligases including Huwe1, COP1, and Pirh2 have all been shown to control the

stability of p53 in cells, underscoring the importance of carefully regulating the levels of this tumour suppressor.² Beyond p53, E3 ubiquitin ligases regulate a number of anti- and pro-apoptotic proteins such as the inhibitor of apoptosis proteins, the Bcl-2 family, and caspases.^{3–7} Given the emerging roles for ubiquitination in development and disease there remains much to be learned about the extent to which E3 ubiquitin ligases regulate apoptosis.

The nematode *Caenorhabditis elegans* is an excellent model organism for studying the role of ubiquitination in apoptosis. This organism has a well-conserved apoptosis signalling pathway that is required for almost all PCDs and a highly conserved complement of E3 ubiquitin ligases.^{8,9} *C. elegans* contains a single p53-like gene, *cep-1*, which is required to activate apoptosis of germ cells treated with DNA-damaging agents.^{10,11} Like mammals, *C. elegans* sense damaged DNA through a conserved checkpoint that includes the 9-1-1 complex (*hpr-9*, *hus-1*, and *mrt-2*) and the Tel2 orthologue *clk-2*. These genes are required to initiate a transient cell-cycle arrest of mitotically proliferating germ cells and/or activate *cep-1*-dependent apoptosis in the pachytene region of the germline.^{12,13} We previously discovered a role for the Skp1/cullin/F-box (SCF) E3 ubiquitin ligase SCF^{FSN-1} in the negative regulation of *cep-1*-dependent germ cell apoptosis.¹⁴ Interestingly, the human equivalent of

¹Developmental and Stem Cell Biology Program, Hospital for Sick Children, Toronto Medical Discovery Tower, 101 College Street, Toronto, Ontario, Canada M5G 1L7 and ²Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada M5S 1A8

*Corresponding author: WB Derry, Developmental and Stem Cell Biology Program, Hospital for Sick Children, MaRS Campus, Room 12-311 TMDT East Tower, 101 College Street, Toronto, Ontario, Canada M5G 1L7. Tel: +416 813 7654, ext. 1829; Fax: +416 813 2212; E-mail: brent.derry@sickkids.ca

Keywords: *C. elegans*; p53 family; apoptosis; DNA damage; E3 ubiquitin ligase; Huwe1/Mule/ARF-BP1

Abbreviations: HECT, homologous to E6-associated protein C terminus; RING, really interesting new gene; PCD, programmed cell death; Mdm2, murine double minute clone 2 oncoprotein; IAP, inhibitor of apoptosis protein; SCF, Skp-Cullin-F-box; IR, ionizing radiation; NGM, nematode growth media; RNAi, RNA interference; HU, hydroxyurea; IPTG, isopropyl- β -D-thiogalactopyranoside

Received 16.3.10; revised 06.12.10; accepted 06.12.10; Edited by JP Medema; published online 14.1.11

SCF^{FSN-1} (SCF^{FBXO45}) was subsequently shown to have a similar role in mammalian cell apoptosis by promoting the proteasome-dependent degradation of the p53 paralogue p73, highlighting the functional conservation of these ligases.¹⁵ The simplified structure of the *C. elegans* cell death pathway and the ease in which genes can be inhibited by RNA interference (RNAi) make this organism an ideal system to study the regulation of DNA damage-induced apoptosis.

To comprehensively examine the role of *C. elegans* 'ubiquitome' in damage-induced apoptosis, we systematically inhibited 108 of the 165 predicted RING, HECT, and U-box-containing E3 ligase genes by RNAi and quantified ionizing radiation (IR)-induced apoptosis in the germline. From this screen, we identified the HECT-domain E3 ligase EEL-1, a homologue of human Huwe1/ARF-BP1/Mule, as a positive regulator of IR-induced germline apoptosis. We show that *eel-1* regulates IR-induced germline apoptosis, but is dispensable for physiological germ cell apoptosis and developmental apoptosis in the soma. EEL-1 promotes apoptosis by a mechanism that does not affect the transcriptional activity of the p53-like protein CEP-1 or alter endogenous levels of the Mcl-1-like protein CED-9. Although *eel-1* mutants exhibit a normal checkpoint response, their progeny are hypersensitive to IR and replication inhibitors. Unexpectedly, the sensitivity of *eel-1* embryos to radiation is not due to an overt defect in DNA repair based on the accumulation of RAD-51 foci in parental germ cells. We conclude that EEL-1 fine-tunes DNA damage-induced apoptosis signalling in the *C. elegans* germline.

Results

The *C. elegans* genome contains a total of 165 predicted E3 ubiquitin ligases, encoding 152 RING proteins, 9 HECT proteins, and 4 U-box proteins.⁹ To determine the extent to which these E3 ligases regulate DNA damage-induced germline apoptosis, we systematically inhibited 108 of these 165 genes by RNAi and quantified germline apoptosis after exposure to 60 Gy of IR. From this screen, we identified *wwp-1* and *eel-1*, two genes encoding HECT-domain containing E3 ligases that caused resistance to IR-induced germline apoptosis when inhibited by RNAi (Figures 1a and b). The *wwp-1* gene has been shown to have a role in nucleotide excision repair in *C. elegans*, and the human homologue WWP1 has been reported to negatively regulate both p53 and p63.^{16–18} However, as we were unable to recapitulate the germline apoptosis phenotype using three mutant alleles of *wwp-1* (*gk372*, *gk397*, and *ok1102*, data not shown) we did not follow up on this gene and instead focused on *eel-1*.

We confirmed our initial *eel-1*(RNAi) observation with two different deletion mutants, which exhibited similar levels of resistance to IR-induced germline apoptosis over a range of dosages (Figures 1c and d). These results suggest that *eel-1* acts a positive regulator of damage-induced apoptosis in the *C. elegans* germline. In the absence of DNA damage, ~50% of germ cells undergo physiological apoptosis.¹⁹ We noticed that physiological germ cell apoptosis was similar in *eel-1* mutants compared with wild-type controls (Figures 1c and d), suggesting that *eel-1* specifically regulates damage-induced




Figure 1 *eel-1* and *wwp-1* are required for promoting IR-induced germ cell apoptosis. (a) Wild-type animals fed *eel-1*(RNAi) (white bars) are resistant to germline apoptosis after treatment with 60 Gy of IR compared with animals fed control RNAi (black bars). (b) *wwp-1*(RNAi) (white bars) also caused resistance to germ cell apoptosis in response to 60 Gy of IR compared with controls. Animals were exposed to IR at the young adult stage and germ cell apoptosis was quantified 24 h later. (c and d) Synchronized L4 hermaphrodites of the indicated genotype were subjected to increasing doses of IR and germ cell corpses were quantified 24 h later. Data represent mean \pm S.E.M. of at least three independent experiments. Between 15 and 30 germlines were scored per experiment

germ cell death. There are two possible explanations for how *eel-1* regulates damage-induced germ cell death. The first is that *eel-1* may control the timing of apoptosis such that germ cell corpses do not appear in *eel-1* mutants until after the 24 h time point used for our analysis. Alternatively, *eel-1* may be required to activate damage-induced germ cell apoptosis similar to the p53-like gene *cep-1*.^{10,11} To test the first possibility, we performed a time course analysis and found that *eel-1(ok1575)* mutants were resistant to IR-induced apoptosis, even 48 h after treatment (Figure 2a). This suggests that *eel-1* is required to promote damage-induced germ cell apoptosis rather than control the kinetics of corpse appearance. As lower levels of IR-induced germ cell apoptosis could be the result of decreased germ cell proliferation in *eel-1* mutants, we determined the number of nuclei in the pachytene region of the germline. The *eel-1* mutants had similar numbers of nuclei in their pachytene regions as wild-type controls (Supplementary Table 1), suggesting that lower levels of apoptosis were not due to a proliferation defect. This is supported by our observation that there are similar numbers of mitotically proliferating germ cells in *eel-1* mutants and wild-type animals (Figures 3a and b). To verify that *eel-1* mutants do not have a general defect in apoptosis, we quantified corpses using the engulfment defective *ced-1(e1735)* allele.²⁰ In *ced-1(e1735); eel-1(ok1575)* double mutants the levels of physiological germline apoptosis

were similar to *ced-1(e1735)* single mutants over the same time course (Figure 2b). To determine if *eel-1* regulates developmental apoptosis in the soma, we compared corpse numbers in developing *ced-1(e1735)* embryos with *ced-1(e1735); eel-1(ok1575)* embryos. Similar to physiological germline apoptosis, we found no difference in the number of persistent corpses in the embryos and first stage larvae of *ced-1; eel-1* versus *ced-1* mutants (Figure 2c). Collectively, these results indicate that *eel-1* specifically regulates DNA damage-induced germ cell apoptosis.

In wild-type animals genotoxic stress activates a conserved DNA damage checkpoint, which causes a transient arrest in mitotically proliferating germ cells and activation of DNA repair. The checkpoint genes *hpr-9/RAD9*, *hus-1/HUS1*, *mrt-2/RAD1*, and *clk-2/TEL2* are required for cell-cycle arrest and activation of DNA repair.^{12,13} Loss-of-function mutations in these genes also cause resistance to germline apoptosis because the checkpoint is unable to activate the p53-like protein CEP-1. As reduced levels of apoptosis in *eel-1* mutants could be the result of defective checkpoint activation, we next examined the checkpoint response in wild-type, *cep-1(gk138)*, and *eel-1(ok1575)* animals. Contrary to van Haaften *et al.*²¹ who reported a mild cell-cycle arrest defect when *eel-1* was knocked down by RNAi, we observed a robust arrest of mitotic germ cell proliferation in *eel-1* deletion mutants that was indistinguishable from wild-type controls




Figure 2 *eel-1* specifically promotes DNA damage-induced germ cell apoptosis. (a) L4 stage hermaphrodites of the indicated genotype were subjected to either 0 or 60 Gy of IR and germ cell corpses were quantified at the indicated times after irradiation. Data represent mean \pm S.E.M. of at least three independent experiments. (b) Physiological germ cell apoptosis is unaffected by mutation of *eel-1*. Germ cell corpses were quantified in strains carrying the *ced-1(e1735)* engulfment defective mutation. Between 15 and 30 germlines were scored per experiment in panels (a) and (b). (c) *eel-1* does not regulate developmental apoptosis. Corpses were quantified at specific stages of embryonic development and first stage (L1) larvae. Around seven animals were scored per stage. Data represent mean \pm S.E.M.




Figure 3 *eel-1* mutants are not defective in DNA damage checkpoint function. (a) *eel-1(ok1575)* mutants undergo a transient arrest in proliferation of mitotic germ cells in response to IR that is indistinguishable from wild-type or *cep-1(gk138)* mutants. Germ lines are outlined in solid black lines and arrows indicate normal mitotic nuclei. Dotted arrows indicate enlarged mitotic nuclei 20 h after exposure to 120 Gy of IR. (b) Quantification of the number of nuclei per unit area in the mitotic region of the germline as shown in panel a. Data represent mean \pm S.E.M. of two independent experiments. Between 10 and 15 germ lines were scored per experiment. (c) Sensitivity of L1 stage *eel-1* mutants to IR-induced sterility is similar to wild type. Twenty animals of the indicated genotype were irradiated at the L1 stage and fertility assessed after they developed into adults

(Figures 3a and b). In addition to defective cell-cycle arrest, checkpoint mutants also generate higher frequencies of inviable progeny after exposure to genotoxic agents because they cannot properly engage DNA repair pathways.^{12,13} Since Huwe1 has recently been shown to regulate DNA repair in mammalian cells,²² we determined if *eel-1* mutants were sensitive to DNA damage by two methods. First, we quantified inviable progeny generated by adults exposed to DNA-damaging agents; second, we determined the frequency of animals irradiated at the L1 stage that developed into sterile adults.²³ We found that *eel-1* mutants generated more inviable progeny than wild-type controls after treatment with IR (Table 1) or replication inhibitors (Supplementary Table 2). In contrast, the percentage of irradiated L1 animals that became sterile by the adult stage was the same as wild type, whereas *clk-2* mutants were hypersensitive (Figure 3c). This apparently contradictory result could be explained if *eel-1*

germlines are not hypersensitive to DNA-damaging agents during larval development but become sensitive by the adult stage.

To address the potential repair function of EEL-1 in more detail, we dissected germlines of irradiated *eel-1* adults and stained with antibodies to the double-strand break repair protein RAD-51. RAD-51 forms foci at sites of double-strand breaks in the germline that increase after treatment with IR and persist for longer periods of time in the germlines of adult checkpoint mutants.^{24,25} We found that the number of persistent RAD-51 foci in *eel-1* mutants was similar to wild-type controls before and after treatment with IR (Figure 4). Interestingly, we observed more RAD-51 foci in irradiated *cep-1; eel-1* double mutants (Figure 4h), suggesting that the combined ablation of these two genes causes defects in DNA repair. However, we cannot exclude the possibility that this could be due to a general metabolic defect in *cep-1; eel-1*

mutants as they also generate more inviable progeny in the absence of genotoxic stress (data not shown). Taken together, the above results suggest that *eel-1* likely acts downstream or independently of the DNA damage checkpoint.

As *cep-1/p53* is required to activate damage-induced germline apoptosis and Huwe1 has been shown to negatively

regulate p53 in mammalian cells, we next asked whether the resistance to IR-induced apoptosis in *eel-1* mutants was the result of decreased CEP-1 transcriptional activity. We previously showed that the E3 ubiquitin ligase SCF^{FSN-1} negatively regulates IR-induced apoptosis through CEP-1, a finding which was subsequently shown to be conserved in humans.^{14,15} To determine if EEL-1 regulates CEP-1 activity, we quantified the transcript abundance of its pro-apoptotic target gene *egl-1* by RT-PCR. Following irradiation, levels of *egl-1* transcript increased by approximately nine-fold in wild-type animals and *eel-1(ok1575)* mutants (Figure 5), suggesting that *eel-1* is not required for CEP-1 activation and likely modulates damage-induced germline apoptosis downstream or independently of *cep-1*.

To determine where *eel-1* acts in relation to the core apoptosis signalling pathway, we examined the epistatic relationship between *eel-1* and the pro-survival gene *ced-9* by constructing a double-mutant strain between the *ced-9(n2812)* putative null allele²⁶ and *eel-1(ok1575)*. As CED-9 functions to protect cells from undergoing apoptosis, loss of

Table 1 The survival of *eel-1* mutants is hypersensitive to treatment with IR

Genotype	Embryonic lethality (%)		
	0 Gy	60 Gy	120 Gy
Wild type	0.5 ± 0.3	30.2 ± 4.0	55.1 ± 7.1
<i>eel-1(ok1575)</i>	12.6 ± 0.1	52.7 ± 3.3	79.3 ± 1.8
<i>eel-1(zu462)</i>	9.0 ± 0.7	57.6 ± 4.1	72.8 ± 1.6
<i>clk-2(qm37)</i>	4.5 ± 1.4	58.7 ± 7.7	81.3 ± 4.9

Synchronized L4 animals were irradiated with the indicated dose and eggs laid 12–18 h later were counted. Unhatched embryos were counted 1 day later to quantify embryonic lethality. Data represent mean ± S.E.M. of three independent experiments




Figure 4 No persistent RAD-51 foci in *eel-1* mutants. RAD-51 foci were scored 24 h after exposure to 30 Gy of IR according to Alpi *et al.*²⁴ Representative images of RAD-51 foci (a–h) and DAPI-stained nuclei (a'–h') are shown for 0 Gy (a–d and a'–d') and after 30 Gy (e–h and e'–h') for wild-type (a, a', e and e'), *eel-1(ok1525)* mutants (b, b', f and f'), *cep-1(gk138)* mutants (c, c', g and g'), and *cep-1(gk138); eel-1(ok1525)* double mutants (d, d', h and h')

CED-9 causes massive germ cell death in adult hermaphrodites. In the absence and presence of damage, homozygous *ced-9(n2812)* hermaphrodites generated from heterozygous mothers had significantly increased levels of germ cell apoptosis compared with wild type, which was not suppressed by the loss of *eel-1* (Figure 6a). This is consistent with a model

where *eel-1* acts upstream or independently of *ced-9*. Complete ablation of *ced-9* in homozygous mutants resulted in similar levels of germline apoptosis in the absence versus presence of IR (Figure 6a). Therefore, if the role of EEL-1 is to specifically regulate CED-9 in response to DNA damage this effect might be masked when the *ced-9* gene is completely ablated. To address this possibility we took two approaches: first, IR-induced germline apoptosis was quantified in a strain heterozygous for the *ced-9(n2812)* allele but homozygous for *eel-1(ok1575)*; second, the levels of endogenous CED-9 protein were examined in *eel-1* mutants by western blotting. Reduction of *ced-9* by one copy in *ced-9(n2812)/hT2* animals resulted in a modest increase in physiological germline apoptosis and hypersensitivity to IR (Figure 6b). In *ced-9(n2812)/hT2; eel-1(ok1575)* animals, IR increased germline apoptosis to levels that were similar to those observed in *ced-9(n2812)/hT2* heterozygotes, suggesting that *eel-1* acts upstream or independently of *ced-9* (Figure 6b). In light of published results that showed Huwe1 to target Mcl-1 (a CED-9 homologue) for degradation in response to DNA damage,⁷ we also quantified the level of CED-9 protein in wild-type animals and *eel-1* mutants before and after DNA damage. There was no significant difference in CED-9 protein levels in *eel-1(ok1575)* or *eel-1(zu462)* mutants compared with wild-type controls following treatment with IR (Figure 6c), supporting a model where EEL-1 does not target CED-9 for degradation in *C. elegans*. Interestingly, we




Figure 5 *eel-1* does not affect the transcriptional activity of the p53-like protein CEP-1. The abundance of the CEP-1 target gene *egl-1* was similar in *eel-1(ok1575)* mutants as wild-type controls, but not induced by IR in *cep-1(gk138)* mutants. Data represent mean \pm S.E.M. of two independent experiments




Figure 6 *eel-1* acts upstream or independently of *ced-9*. (a and b) Hermaphrodites of the indicated genotypes were irradiated with 60 Gy of IR at the L4 larval stage and germ cell apoptosis quantified 24 h later. Data represent mean \pm S.E.M. of three independent experiments. Between 15 and 30 germlines were scored per experiment. (c) Endogenous CED-9 protein levels are not affected by *eel-1* status or IR. Hermaphrodites of the indicated genotype were treated with 120 Gy of IR at the L4 stage and the protein was extracted from whole-worm lysates after 24 h. Western analysis was performed using an anti-CED-9 antibody kindly provided by Dr. Barbara Conratt. The protein α tubulin was used as a loading control. Quantification was performed using ImageJ software. Protein abundance is relative to wild type at the indicated dose of IR

found CED-9 protein levels to be unchanged in unirradiated wild-type animals compared with irradiated worms, suggesting that CED-9 stability does not change in response to DNA damage. However, it is possible that CED-9 protein is regulated by DNA damage only in the pachytene region of the germline, which might not be detectable by western analysis from whole-worm lysates.

As *eel-1* appears to function downstream of *cep-1* and does not regulate CED-9 protein levels, we sought to identify other possible substrates of EEL-1. Previous work revealed a role for EEL-1 in cell fate specification via the transcription factor SKN-1, which regulates mesendodermal cell fate specification in the early embryo and is the only proposed substrate of EEL-1 in *C. elegans*.²⁷ As EEL-1 and SKN-1 have been proposed to interact in the early embryo to regulate development, it is possible that they may also act in the same pathway to regulate apoptosis in the germline. If SKN-1 is a target of EEL-1 in germ cell apoptosis signalling, ablation of *skn-1* might increase sensitivity to DNA damage-induced germ cell apoptosis. While ablation of *skn-1* by RNAi and a loss-of-function mutation in *skn-1* caused the expected embryonic lethality in the next generation, IR-induced germ cell apoptosis was indistinguishable from wild-type controls (Supplementary Figure 1, and data not shown). These results suggest that *skn-1* does not have a major role in DNA damage-induced germ cell apoptosis and that EEL-1 likely targets a novel substrate.

Discussion

We screened 65% of the 165 predicted E3 ligase genes comprising the HECT, RING, and U-box domain subgroups and identified a pro-apoptotic role for *eel-1*. We also uncovered the WWP1 homologue (*wwp-1*) in this screen and found a similar pro-apoptotic role as *eel-1* in germline apoptosis. However, as existing deletion alleles of *wwp-1* did not recapitulate the apoptotic phenotypes observed by RNAi we focused on characterizing *eel-1*. The mammalian homologue of EEL-1, Huwe1 (also known as ARF-BP1, HectH9, or Mule) has been shown to regulate both pro-apoptotic (p53) and anti-apoptotic (Mcl-1) proteins, as well as the DNA repair proteins Cdc6 and DNA polymerase β .^{7,22,28,29} Our results suggest that EEL-1 regulates apoptosis in *C. elegans* by a different mechanism than its mammalian counterpart. As there is a robust cell-cycle arrest and the pro-apoptotic gene *egl-1* is induced to wild-type levels in *eel-1* mutants exposed to IR (Figures 3 and 5), our results imply that *eel-1* functions in a parallel signalling pathway to promote damage-induced germline apoptosis. Consistent with this, several recent reports have uncovered genes that act independently or downstream of *cep-1* to promote apoptosis, such as the Sirtuin *sir-2.1*, the Rb gene *lin-35*, the inhibitor of growth homologue *ing-3* and the Krit1/Ccm1 gene *kri-1*.^{30–33} Like *eel-1*, all of these genes have pro-apoptotic function independent of the DNA damage checkpoint or CEP-1 transcriptional activity. In *C. elegans*, EEL-1 has been shown to bind and possibly stimulate degradation of the transcription factor SKN-1 in the early embryo.²⁷ We did not observe any changes in IR-induced germline apoptosis after knocking down *skn-1* by RNAi or in *skn-1* mutants (Supplementary Figure 1),

suggesting that EEL-1 targets a novel substrate to regulate apoptosis.

The hypersensitivity of *eel-1* mutants to IR-induced lethality (Table 1) is intriguing in light of a recent study implicating Huwe1 in the regulation of base excision repair by targeting DNA polymerase β .²² Interestingly, *clk-2* checkpoint mutants are also hypersensitive to IR and replication stress and exhibit synthetic lethal interactions with *cep-1*, suggesting that *cep-1* cooperates with *clk-2* to modulate replication stress during embryonic development.^{12,34} However, there was no difference in the accumulation and persistence of RAD-51 foci in the germlines of *eel-1* mutants compared with wild-type controls exposed to IR (Figure 4). We conclude that either EEL-1 has no role in DNA repair or it regulates repair independent of RAD-51 accumulation. Interestingly, *cep-1*; *eel-1* double mutants generate about 40% lethal progeny in the absence of exogenous DNA damage, similar to *cep-1*; *clk-2* double mutants,³⁴ and they appear to have more intense RAD-51 foci after treatment with IR (Figure 4). Thus, it is possible that these genes have a cooperative role in DNA repair. The hypersensitivity of *eel-1* mutants to the replication inhibitor hydroxyurea (HU) (Supplementary Table 2) provides additional evidence that this E3 ligase may target a unique substrate downstream of the checkpoint to coordinate repair and apoptosis. However, strains carrying either allele of *eel-1* exhibit a range of pleiotropic phenotypes, such as a protruding vulva and egg laying defects. Thus, it is possible that *eel-1* mutants are hypersensitive to stresses that simply antagonize an already tenuous existence. As Huwe1 has been shown to target multiple proteins that regulate apoptosis and DNA repair in mammals, it is possible that EEL-1 also targets multiple substrates in *C. elegans*. Consistent with this, EEL-1 has been implicated in the control of cell fate specification through SKN-1 and more recently in hemidesmosome maturation in *C. elegans*.^{27,35} Therefore, ablation of the *eel-1* gene likely deregulates several cellular proteins and signalling pathways that could indirectly cause animals to become hypersensitive to stresses such as DNA damage. More work will have to be done to determine whether EEL-1 has a direct role in DNA repair. However, we believe that *eel-1* has a specific role in the regulation of DNA damage-induced apoptosis as there are no proliferation defects, no changes in physiological germline apoptosis, and no effects on developmental apoptosis in *eel-1* mutants. As IR-induced apoptosis in *eel-1*; *ced-9* double mutants is the same as *ced-9* single mutants (Figure 6) we conclude that EEL-1 does not likely affect the core apoptotic pathway downstream of *egl-1*. This is supported by the absence of defects in physiological germline apoptosis or developmental apoptosis in *eel-1* mutants (Figure 2).

How might *eel-1* regulate DNA damage-induced germline apoptosis? Taken together, our data suggest that *eel-1* regulates apoptosis downstream or independently of *cep-1* and the DNA damage checkpoint (Figure 7). It is unlikely that EEL-1 targets CED-9 directly, as changes in endogenous CED-9 levels (or mobility shifts) were not observed in *eel-1* mutants following treatment with IR (Figure 6). However, as CED-9 is expressed in most cells of the organism it is possible that alterations in the pachytene region of the germline might not be detectable by western analysis of whole worm lysates.




Figure 7 Model for EEL-1 regulation of IR-induced germ cell apoptosis in *C. elegans*. EEL-1 may act at the level of EGL-1 or CED-9. Hypothetical relationships are indicated by dashed arrows. The corresponding mammalian pathway and the EEL-1 orthologue Huwe1 are shown on the right

In general, polyubiquitin chains generated through linkages at residues K11, K24, or K48 cause substrates to be directed for degradation by the proteasome. On the other hand, mono-ubiquitination or polyubiquitination via linkage at K63 can alter the activity and/or subcellular localization of proteins.¹ As the BH3-only gene *egl-1* is also dispensable for physiological germ cell apoptosis,¹⁹ it is possible that EEL-1 could monoubiquitinate EGL-1 (or an EGL-1 accessory protein) to control its activity and/or subcellular localization in response to DNA damage. It is well established that *egl-1* is upregulated transcriptionally to promote apoptosis during development and in the germline in response to genotoxic stress.^{8,36,37} However, it is not known whether EGL-1 is regulated by post-translational modifications in *C. elegans*. Post-translational modification by ubiquitin has been shown to have a role in the regulation of BH3-only proteins in mammalian cells. For example, the BH3-only protein BIM, which is responsible for the apoptosis of lymphocytes, myeloid cells, and neurons, is ubiquitinated and degraded in human osteoclasts by the c-Cbl family of E3 ubiquitin ligases.³⁸ It will be important to determine whether this aspect of BH3-only protein regulation is conserved in *C. elegans*, which can be assessed once antibodies to EGL-1 become available. Although further studies are required to elucidate the biochemical mechanism by which EEL-1 regulates damage-induced germline apoptosis, our results suggest that this ligase may have an ancient role in apoptosis signalling that operates by a distinct mechanism in *C. elegans*.

Materials and Methods

General methods and strains. All *C. elegans* strains used were cultured at 20°C on nematode growth media (NGM) plates according to standard protocols.³⁹ The wild-type strain used was Bristol N2. The following strains were provided by the *Caenorhabditis* Genetics Centre (CGC): *ced-1(e1735)*, *cep-1(gk138)*, *wpp-1(gk372)*, *wpp-1(ok1102)*, *wpp-1(gk397)*, *dpy-5(e61) unc-13(e51)/hT2[bli-4(e937) let-?(q782) qIs48]*, *hus-1(op244) l/hT2[bli-4(e937) let-?(q782) qIs48]* (l;III), *clk-2(qm37)*, *eel-1(ok1575)*, and *eel-1(zu462) unc-33(e204)*. The strain *ced-9(n2812)/qC1 III* was kindly provided by Dr. Robert Horvitz. The strain

ced-9(n2812)/hT2 III was generated with *dpy-5(e61) unc-13(e51)/hT2[bli-4(e937) let-?(q782) qIs48]* and *ced-9(n2812)/qC1 III* using standard genetic techniques. The double mutants *ced-9(n2812)/hT2 III; eel-1(ok1575)* and *ced-1(e1735); eel-1(ok1575)* were generated using standard methods. The linked *unc-33(e204)* mutation was segregated away from *eel-1(zu462) unc-33(e204)* to generate *eel-1(zu462)*.

Screening and RNAi. RNAi experiments were performed by the feeding method and all of the RNAi bacterial strains were obtained from the Ahinger RNAi library as we previously described.¹⁴ A bacterial strain expressing the empty RNAi vector L4440 was used as a control. To screen the 108 E3 ligase genes represented in the Ahinger RNAi library for defects in IR-induced germline apoptosis, wild-type animals of the fourth larval stage (L4) were fed double-stranded RNA expressing bacteria and were allowed to lay eggs for 3–4 days. Young adult animals of the F1 generation were treated with 60 Gy of IR and germline apoptosis was quantified as described (see below). All 108 E3 ligases available in the RNAi library were screened at least once.

***C. elegans* germline apoptosis.** Synchronized hermaphrodites at the L4 stage of development were exposed to various doses of IR (¹³⁷Cs source) and were allowed to recover on NGM plates at 20°C. Germ cell apoptosis was quantified in the pachytene region of one gonad arm per animal using Nomarski optics 24 h post-irradiation as previously described.¹⁰ As older *eel-1* mutants exhibit Pvl and Egl phenotypes, we scored germline apoptosis in worms 24 h after the L4 molt. Developmental apoptosis in staged embryos and physiological germ cell apoptosis in young adult hermaphrodites were quantified in strains carrying the *ced-1(e1735)* mutation to prevent corpse engulfment.¹⁹

Cell-cycle arrest and radiation sensitivity. To determine the effects of IR on germ cell proliferation, synchronized L4 animals were treated with 120 Gy of IR and cell nuclei per area in the mitotic region of the germline were quantified 24 h after irradiation. Analysis was performed at $\times 630$ using Nomarski microscopy and OpenLab software (PerkinElmer, Waltham, MA, USA), as previously described.⁴⁰

To evaluate radiation sensitivity, L4-stage hermaphrodites were treated with either 60 or 120 Gy of IR. Approximately 12 h later, irradiated worms and unirradiated controls were transferred to new plates and allowed to lay eggs for 5 h (eggs laid during this time correspond to germ cells irradiated at pachytene). The adults were removed and the number of unhatched eggs were counted 24 h later. Radiation-induced sterility was evaluated by exposing L1 stage hermaphrodites to IR and quantifying the percentage of animals that developed into sterile adults.²³

Characterization of *eel-1* mutants. To determine if *ok1575* and *zu462* are recessive alleles, we crossed these *eel-1* mutants with wild-type animals and quantified damage-induced germ cell apoptosis in the heterozygous progeny. The *eel-1* heterozygous mutants had similar levels of germ cell apoptosis as wild-type controls treated with 60 Gy of IR (data not shown), indicating that *ok1575* and *zu462* are recessive alleles.

Both *eel-1(ok1575)* and *eel-1(zu462)* mutants also display some pleiotropic defects such as protruding vulvas, slow growth, and egg-laying defects. It is unknown whether these are null alleles, as no deficiencies are available that remove the *eel-1* locus. The *eel-1(ok1575)* allele contains a 1.6-kb in-frame deletion and RT-PCR analysis revealed that a truncated *eel-1* transcript is expressed in these animals (data not shown). The *eel-1(zu462)* mutant isolated by Page et al.²⁷ contains a 1.9-kb deletion followed by a 5.9-kb inverted duplication that is predicted to encode an extra 36 amino acids and generate a truncated protein of 3911 amino acids.

***egl-1* quantitative real-time PCR.** Approximately 2000 synchronized L4 stage animals were treated with either 0 or 120 Gy of IR and RNA was isolated 24 h later. Worms were washed twice in M9 buffer and stored at –80°C overnight in 1 ml of Trizol (Invitrogen Canada, Burlington, ON, Canada). The next day lysates were thawed and RNA was extracted with 0.2 ml of chloroform then centrifuged at 15 000 r.p.m. for 15 min at 4°C. To precipitate RNA, 200 μ l of supernatant was placed in a fresh 1.5-ml microcentrifuge tube plus 500 μ l of isopropanol then centrifuged at 15 000 r.p.m. for 15 min. The pellet was washed with 1 ml of 75% ethanol in DEPC H₂O (Invitrogen) and centrifuged at 15 000 r.p.m. for 5 min. The pellet was air dried for 15 min then re-suspended in 25 μ l DEPC H₂O.

A cDNA template was generated using 2 μ g of RNA, as previously described.³⁹ The *egl-1* transcript was amplified using primers described by Hofmann et al.³⁶

Amplification of tubulin transcript was used as a control. Real-time monitoring of amplification was achieved using a 7900HT real-time PCR system (Applied Biosystems, Foster City, CA, USA).

Western blot analysis of CED-9 protein levels. Approximately 2000–3000 synchronized hermaphrodites at the L4 stage of development were exposed to either 0 or 120 Gy of IR. Animals were lysed by sonication in SDS lysis buffer (4% SDS, 100 mM Tris-HCl (pH 6.8), 20% glycerol) 24 h after treatment with IR. In total, 30 µg of soluble protein was suspended in SDS loading buffer (4% SDS, 10% 2-mercaptoethanol, 20% glycerol, 0.004% bromophenol blue, 100 mM Tris-HCl (pH 6.8)) and was boiled for 4 min and then centrifuged at 12 000 × g for 1 min. Proteins were separated by SDS-PAGE, transferred onto a nitrocellulose membrane, and immunoblotted with a 1/1000 dilution of rabbit anti-CED-9 antibody (kindly provided by Dr. Barbara Conradt) or a 1/2000 dilution of mouse anti- α -tubulin antibody (Sigma-Aldrich Canada Ltd, Oakville, ON, Canada). ImageJ software (<http://rsbweb.nih.gov/ij/index.html>) was used to analyse bands corresponding to CED-9 and α -tubulin and quantification was performed by normalizing CED-9 protein levels to α -tubulin protein levels.

Immunocytochemistry. Worms at the young adult stage were exposed to either 0 or 30 Gy of IR and allowed to recover for 24 h at 20°C. Irradiated or unirradiated control hermaphrodites were washed once in PBS, resuspended in 30 µl of PBS + 0.2 mM tetramisole on polylysine-coated slides. Germlines were extruded by removing the heads and tails using scissoring motion with two 27-G needles. Tetramisole was replaced with 1% paraformaldehyde in PBS for 10 min at room temperature followed by freeze cracking on dry ice for 1 min. Post fixation was carried by immersing samples in a 1:1 mixture of methanol/acetone at –20°C for 1 min. Samples were then permeabilized with PBS + 1% Triton X-100 for 10 min at room temperature, then washed three times with PBS + 0.1% Tween 20 (PBST) for 10 min per wash. To reduce background, samples were incubated with Image-IT FX signal enhancer (Invitrogen) for 20 min, then incubated for 30 min in PBS + 0.1% Tween 20 + 1% BSA (PBSTB). RAD-51 antibodies (SDI, cat no. 2948.00.02) were diluted 1/100 in PBSTB and incubated at 4°C overnight in a humid chamber. Samples were washed three times in PBST then samples were incubated in Alexa 488 (Invitrogen) secondary antibody diluted 1/1000 in PBSTB for 1 h at room temperature. After washing four times for 15 min in PBST, the samples were mounted in prolong Gold (Invitrogen). DAPI diluted 1/1000 from 1 mg/ml stock solution was added at the second wash step.

HU sensitivity. NGM plates were prepared with 10 mM HU in the media as described previously.³⁴ Hermaphrodites at the L4 stage were placed on HU supplemented NGM plates and allowed to develop into young adults for 24 h then transferred to fresh HU plates and allowed to lay eggs for 6 h. The adults were then removed and the deposited eggs were counted. Unhatched eggs were counted after 48 h and the total number of living larvae and adult animals were counted after 96 h. Percent lethality was determined in triplicate.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements. We thank the *Caenorhabditis* Genetics Center, which is funded by the National Institutes of Health Center for Research Resources for providing worm strains. We kindly thank H.R. Horvitz (Massachusetts Institute of Technology) for providing the *ced-9(n2812)/qC1* strain, B. Conradt and Stephane Rolland (Dartmouth University) for CED-9 antibody and assistance with the western blot protocol. This work was supported by a Canadian Institutes of Health Research award to W.B.D., Natural Sciences and Engineering Research Council of Canada Graduate Scholarship to A.R., and a Canadian Institutes of Health Strategic Training Fellow in the Excellence in Radiation Research for the 21st Century Program to M.L. We are grateful to A. Perrin and S. Ito for helpful input on the manuscript.

- Pickart CM. Mechanisms underlying ubiquitination. *Annu Rev Biochem* 2001; **70**: 503–533.
- Brooks CL, Gu W. p53 ubiquitination: Mdm2 and beyond. *Mol Cell* 2006; **21**: 307–315.

- Dimmeler S, Breitschopf K, Haendeler J, Zeiher AM. Dephosphorylation targets Bcl-2 for ubiquitin-dependent degradation: a link between the apoptosome and the proteasome pathway. *J Exp Med* 1999; **189**: 1815–1822.
- Li B, Dou QP. Bax degradation by the ubiquitin/proteasome-dependent pathway: involvement in tumor survival and progression. *Proc Natl Acad Sci USA* 2000; **97**: 3850–3855.
- Marshansky V, Wang X, Bertrand R, Luo H, Duguid W, Chinnadurai G et al. Proteasomes modulate balance among proapoptotic and antiapoptotic Bcl-2 family members and compromise functioning of the electron transport chain in leukemic cells. *J Immunol* 2001; **166**: 3130–3142.
- Yang Y, Fang S, Jensen JP, Weissman AM, Ashwell JD. Ubiquitin protein ligase activity of IAPs and their degradation in proteasomes in response to apoptotic stimuli. *Science* 2000; **288**: 874–877.
- Zhong Q, Gao W, Du F, Wang X. Mule/ARF-BP1, a BH3-only E3 ubiquitin ligase, catalyzes the polyubiquitination of Mcl-1 and regulates apoptosis. *Cell* 2005; **121**: 1085–1095.
- Conradt B, Xue D. Programmed cell death. *WormBook* 2005; **6**: 1–13.
- Kipreos ET. Ubiquitin-mediated pathways in *C. elegans*. *WormBook* 2005: 1–24.
- Derry WB, Putzke AP, Rothman JH. *Caenorhabditis elegans* p53: role in apoptosis, meiosis, and stress resistance. *Science* 2001; **294**: 591–595.
- Schumacher B, Hofmann K, Boulton S, Gartner A. The *C. elegans* homolog of the p53 tumor suppressor is required for DNA damage-induced apoptosis. *Curr Biol* 2001; **11**: 1722–1727.
- Ahmed S, Alpi A, Hengartner MO, Gartner A. *C. elegans* RAD-51/CLK-2 defines a new DNA damage checkpoint protein. *Curr Biol* 2001; **11**: 1934–1944.
- Gartner A, Milstein S, Ahmed S, Hodgkin J, Hengartner MO. A conserved checkpoint pathway mediates DNA damage-induced apoptosis and cell cycle arrest in *C. elegans*. *Mol Cell* 2000; **5**: 435–443.
- Gao MX, Liao EH, Yu B, Wang Y, Zhen M, Derry WB. The SCF^{FSN-1} ubiquitin ligase controls germline apoptosis through CEP-1/p53 in *C. elegans*. *Cell Death Differ* 2008; **15**: 1054–1062.
- Peschiarioli A, Scialpi F, Bernassola F, Pagano M, Melino G. The F-box protein FBXO45 promotes the proteasome-dependent degradation of p73. *Oncogene* 2009; **28**: 3157–3166.
- Astin JW, O'Neil NJ, Kuwabara PE. Nucleotide excision repair and the degradation of RNA pol II by the *Caenorhabditis elegans* XPA and Rsp5 orthologues, RAD-3 and WWP-1. *DNA Repair (Amst)* 2008; **7**: 267–280.
- Laine A, Ronai Z. Regulation of p53 localization and transcription by the HECT domain E3 ligase WWP1. *Oncogene* 2007; **26**: 1477–1483.
- Li Y, Zhou Z, Chen C. WW domain-containing E3 ubiquitin protein ligase 1 targets p63 transcription factor for ubiquitin-mediated proteasomal degradation and regulates apoptosis. *Cell Death Differ* 2008; **15**: 1941–1951.
- Gumienny TL, Lambie E, Hartwig E, Horvitz HR, Hengartner MO. Genetic control of programmed cell death in the *Caenorhabditis elegans* hermaphrodite germline. *Development* 1999; **126**: 1011–1022.
- Hedgecock EM, Sulston JE, Thomson JN. Mutations affecting programmed cell deaths in the nematode *Caenorhabditis elegans*. *Science* 1983; **220**: 1277–1279.
- van Haaften G, Romeijn R, Pothof J, Koole W, Mullenders LH, Pastink A et al. Identification of conserved pathways of DNA-damage response and radiation protection by genome-wide RNAi. *Curr Biol* 2006; **16**: 1344–1350.
- Parsons JL, Tait PS, Finch D, Dianova II, Edelmann MJ, Khoronenkova SV et al. Ubiquitin ligase ARF-BP1/Mule modulates base excision repair. *EMBO J* 2009; **28**: 3207–3215.
- Bailly AP, Freeman A, Hall J, Declais AC, Alpi A, Lilley DM et al. The *Caenorhabditis elegans* homolog of Gen1/Yen1 resolves links DNA damage signaling to DNA double-strand break repair. *PLoS Genet* 2010; **6**: e1001025.
- Alpi A, Pasierbek P, Gartner A, Loidl J. Genetic and cytological characterization of the recombination protein RAD-51 in *Caenorhabditis elegans*. *Chromosoma* 2003; **112**: 6–16.
- Martin JS, Winkelmann N, Petalcorin MI, McIlwraith MJ, Boulton SJ. RAD-51-dependent and -independent roles of a *Caenorhabditis elegans* BRCA2-related protein during DNA double-strand break repair. *Mol Cell Biol* 2005; **25**: 3127–3139.
- Hengartner MO, Horvitz HR. *C. elegans* cell survival gene *ced-9* encodes a functional homolog of the mammalian proto-oncogene bcl-2. *Cell* 1994; **76**: 665–676.
- Page BD, Diede SJ, Tenlen JR, Ferguson EL. EEL-1, a Hect E3 ubiquitin ligase, controls asymmetry and persistence of the SKN-1 transcription factor in the early *C. elegans* embryo. *Development* 2007; **134**: 2303–2314.
- Chen D, Kon N, Li M, Zhang W, Qin J, Gu W. ARF-BP1/Mule is a critical mediator of the ARF tumor suppressor. *Cell* 2005; **121**: 1071–1083.
- Hall JR, Kow E, Nevis KR, Lu CK, Luce KS, Zhong Q et al. Cdc6 stability is regulated by the Huw1 ubiquitin ligase after DNA damage. *Mol Biol Cell* 2007; **18**: 3340–3350.
- Greiss S, Hall J, Ahmed S, Gartner A. *C. elegans* SIR-2.1 translocation is linked to a proapoptotic pathway parallel to *cep-1/p53* during DNA damage-induced apoptosis. *Genes Dev* 2008; **22**: 2831–2842.
- Ito S, Greiss S, Gartner A, Derry WB. Cell-nonautonomous regulation of *C. elegans* germ cell death by *kri-1*. *Curr Biol* 2005; **15**: 333–338.

32. Luo J, Shah S, Riabowol K, Mains PE. The *Caenorhabditis elegans* *ing-3* gene regulates ionizing radiation-induced germ-cell apoptosis in a p53-associated pathway. *Genetics* 2009; **181**: 473–482.
33. Schertel C, Conradt B. *C. elegans* orthologs of components of the RB tumor suppressor complex have distinct pro-apoptotic functions. *Development* 2007; **134**: 3691–3701.
34. Derry WB, Bierings R, van Iersel M, Satkunendran T, Reinke V, Rothman JH. Regulation of developmental rate and germ cell proliferation in *Caenorhabditis elegans* by the p53 gene network. *Cell Death Differ* 2007; **14**: 662–670.
35. Zahreddine H, Zhang H, Diogon M, Nagamatsu Y, Labouesse M. CRT-1/calreticulin and the E3 ligase EEL-1/HUWE1 control hemidesmosome maturation in *C. elegans* development. *Curr Biol* 2010; **20**: 322–327.
36. Hofmann ER, Milstein S, Boulton SJ, Ye M, Hofmann JJ, Stergiou L *et al*. *Caenorhabditis elegans* HUS-1 is a DNA damage checkpoint protein required for genome stability and EGL-1-mediated apoptosis. *Curr Biol* 2002; **12**: 1908–1918.
37. Nehme R, Conradt B. *egl-1*: a key activator of apoptotic cell death in *C. elegans*. *Oncogene* 2008; **27** (Suppl 1): S30–S40.
38. Akiyama T, Bouillet P, Miyazaki T, Kadono Y, Chikuda H, Chung UI *et al*. Regulation of osteoclast apoptosis by ubiquitylation of proapoptotic BH3-only Bcl-2 family member Bim. *EMBO J* 2003; **22**: 6653–6664.
39. Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics* 1974; **77**: 71–94.
40. Quevedo C, Kaplan DR, Derry WB. AKT-1 regulates DNA-damage-induced germline apoptosis in *C. elegans*. *Curr Biol* 2007; **17**: 286–292.

Supplementary Information accompanies the paper on Cell Death and Differentiation website (<http://www.nature.com/cdd>)